

Measurement of Branching Fractions and Charge Asymmetries for Exclusive B Decays to Charmonium

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges-Pous,² A. Palano,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ A. W. Borgland,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. E. Latham,⁹ F. F. Wilson,⁹ T. Cuhadar-Donszelmann,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ D. Thiessen,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² V. N. Ivanchenko,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfiel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ Sh. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ A. Lu,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzki,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ S. Yang,¹⁹ S. Jayatilleke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ J. Zhang,²¹ L. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² J. L. Harton,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² B. Spaan,²³ D. Altenburg,²⁴ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ E. Feltresi,²⁴ A. Hauke,²⁴ H. M. Lacker,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ A. Sarti,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ G. Crosetti,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² M. J. Charles,³³ G. J. Grenier,³³ U. Mallik,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ J. Lamsa,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ T. C. Petersen,³⁵ S. Plaszczynski,³⁵ M. H. Schune,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹ R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ M. A. Winter,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³ V. B. Koptchev,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸

P. Taras,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,*} F. Fabozzi,^{50,*} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ Ch. de la Vaissière,⁵⁶ L. Del Buono,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malcles,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ M. Bondioli,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ G. Simi,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ V. Miftakov,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,^{61,62} A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzoni,⁶² S. Morganti,⁶² M. Pierini,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Schott,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ G. De Nardo,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ J. Strube,^{54,67} D. Su,⁶⁷ M. K. Sullivan,⁶⁷ J. Thompson,⁶⁷ J. Va'vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ H. Kim,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ L. Bosio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,^{74,†} L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ F. Martinez-Vidal,⁷⁵ R. S. Panvini,⁷⁶ Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ J. Wu,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ M. G. Greene,⁸⁰ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

- ¹⁹California Institute of Technology, Pasadena, California 91125, USA
- ²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA
- ²¹University of Colorado, Boulder, Colorado 80309, USA
- ²²Colorado State University, Fort Collins, Colorado 80523, USA
- ²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
- ²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
- ²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
- ²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- ²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ³⁰Harvard University, Cambridge, Massachusetts 02138, USA
- ³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³²Imperial College London, London, SW7 2AZ, United Kingdom
- ³³University of Iowa, Iowa City, Iowa 52242, USA
- ³⁴Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁵Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France
- ³⁶Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁷University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ³⁸Queen Mary, University of London, E1 4NS, United Kingdom
- ³⁹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ⁴⁰University of Louisville, Louisville, Kentucky 40292, USA
- ⁴¹University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴²University of Maryland, College Park, Maryland 20742, USA
- ⁴³University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴⁴Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁵McGill University, Montréal, Quebec, Canada H3A 2T8
- ⁴⁶Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁷University of Mississippi, University, Mississippi 38677, USA
- ⁴⁸Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7
- ⁴⁹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁵⁰Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵¹NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵²University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵³Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁴University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁵Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁶Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
- ⁵⁷University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁵⁸Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ⁵⁹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶⁰Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶¹Princeton University, Princeton, New Jersey 08544, USA
- ⁶²Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶³Universität Rostock, D-18051 Rostock, Germany
- ⁶⁴Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁵DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁶University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁷Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁶⁸Stanford University, Stanford, California 94305-4060, USA
- ⁶⁹State University of New York, Albany, New York 12222, USA
- ⁷⁰University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷¹University of Texas at Austin, Austin, Texas 78712, USA
- ⁷²University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷³Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁴Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁵Universidad de Valencia, E-46100 Burjassot, Valencia, Spain
- ⁷⁶Vanderbilt University, Nashville, Tennessee 37235, USA
- ⁷⁷University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁸Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
- ⁷⁹University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁸⁰Yale University, New Haven, Connecticut 06511, USA

(Dated: February 7, 2008)

We report measurements of branching fractions and charge asymmetries of exclusive decays of neutral and charged B mesons into two-body final states containing a charmonium state and a light strange meson. The charmonium mesons considered are J/ψ , $\psi(2S)$ and χ_{c1} , and the light meson is either K or K^* . We use a sample of about 124 million $B\bar{B}$ pairs collected with the BABAR detector at the PEP-II storage ring at the Stanford Linear Accelerator Center.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

Nonleptonic decays of B mesons provide tests of both strong- and weak-interaction dynamics. Decays $B \rightarrow (c\bar{c})K^{(*)}$ are particularly illuminating as they involve three kinds of mesons: one with a heavy quark and a light quark, one with two heavy quarks, and one with two light quarks. Phenomenological models (see Refs.[2-12] of Ref. [1]) give estimates for branching fractions and for ratios of the decays to K and K^* . Some branching fractions and ratios have been reported but others have not, so more stringent tests of the models are possible. The Standard Model predicts small differences between the branching fractions for positive and negative B mesons, i.e. small direct CP violation [2]. Large charge asymmetries would be evidence for new physics. Limits on direct CP violation would constrain extensions of the Standard Model. Very few measurements have been reported in $B \rightarrow (c\bar{c})K^{(*)}$ modes [3]. The decay processes studied in this paper are listed in Table I.

TABLE I: Decay processes studied in this analysis. The $(c\bar{c})$ resonance is either a J/ψ , $\psi(2S)$, or χ_{c1} . For all processes, when relevant, we use the secondary decay modes $J/\psi, \psi(2S) \rightarrow \ell^+\ell^-$, $\chi_{c1} \rightarrow J/\psi \gamma$, and $K_S^0 \rightarrow \pi^+\pi^-$, $\pi^0 \rightarrow \gamma\gamma$.

Decay Channel	Secondary K^* decay mode
$B^0 \rightarrow (c\bar{c}) K^{*0}$	$K^{*0} \rightarrow K^+ \pi^-, K_S^0 \pi^0$
$B^+ \rightarrow (c\bar{c}) K^{*+}$	$K^{*+} \rightarrow K^+ \pi^0, K_S^0 \pi^+$
$B^0 \rightarrow (c\bar{c}) K_S^0$	
$B^+ \rightarrow (c\bar{c}) K^+$	

The data sample used in this analysis contains 124 million $B\bar{B}$ events collected with the BABAR detector at the PEP-II asymmetric e^+e^- storage ring. This represents a total integrated luminosity of 112.4 fb^{-1} taken on the $\Upsilon(4S)$ resonance. The BABAR detector is described in detail elsewhere [4]. Surrounding the interaction point, a five-layer double-sided silicon vertex tracker (SVT) provides precise reconstruction of track angles and B decay vertices. A 40-layer drift chamber (DCH) provides measurements of the transverse momenta of charged particles. An internally reflecting ring-imaging Cherenkov detector (DIRC) is used for particle identification. A CsI(Tl) crystal electromagnetic calorimeter (EMC) is used to detect photons and electrons. The calorimeter is surrounded by a 1.5-T magnetic field. The flux return is instrumented with resistive plate chambers (RPC) used for muon and neutral-hadron identification.

Multihadron events are selected by demanding a minimum of three reconstructed charged tracks in the polar-angle range $0.41 < \theta < 2.54 \text{ rad}$, where θ is defined in the laboratory frame. Charged tracks must be reconstructed in the DCH and are required to originate within 1.5 cm of the beam in the plane transverse to it and within 10 cm of the beamspot along the beam direction. Events are required to have a primary vertex within 0.5 cm of the average position of the interaction point in the plane transverse to the beamline, and within 6 cm longitudinally. Charged tracks are required to include at least 12 DCH hits and to have a transverse momentum $p_T > 100 \text{ MeV}/c$. Photons are reconstructed from EMC clusters. The lateral energy profile (LAT) [5] is used to discriminate electromagnetic from hadronic clusters. Photons are required to have a minimum energy of 30 MeV, to satisfy $\text{LAT} < 0.8$, and to be in the fiducial volume $0.41 < \theta < 2.41 \text{ rad}$. Electron candidates are selected using information from the EMC (LAT and Zernike moment A_{42} [6]), the ratio of the energy measured in the EMC to the momentum measured by the tracking system, the energy loss in the drift chamber, and the Cherenkov angle measured in the DIRC. Electrons are also required to be in the fiducial volume $0.41 < \theta < 2.41 \text{ rad}$. Muon candidates are selected using information from the EMC (energy deposition consistent with a minimum ionizing particle) and the distribution of hits in the RPC. Muons are required to be in the fiducial volume $0.3 < \theta < 2.7 \text{ rad}$. We select charged kaon and pion candidates using information from the energy loss in the SVT and DCH, and the Cherenkov angle measured in the DIRC. Kaon candidates are required to be in the fiducial volume $0.45 < \theta < 2.45 \text{ rad}$.

The selection has been optimized by maximizing the ratio $S/\sqrt{S+B}$, where S and B are the number of expected signal and background events obtained from Monte Carlo simulation. The J/ψ candidates are required to have an invariant mass $2.95 < M_{e^+e^-} < 3.14 \text{ GeV}/c^2$ or $3.06 < M_{\mu^+\mu^-} < 3.14 \text{ GeV}/c^2$ for $J/\psi \rightarrow e^+e^-$ or $J/\psi \rightarrow \mu^+\mu^-$ decays respectively. The $\psi(2S)$ candidates are required to have an invariant mass $3.44 < M_{e^+e^-} < 3.74 \text{ GeV}/c^2$ or $3.64 < M_{\mu^+\mu^-} < 3.74 \text{ GeV}/c^2$ for $\psi(2S) \rightarrow e^+e^-$ or $\psi(2S) \rightarrow \mu^+\mu^-$ decays respectively. For $J/\psi, \psi(2S) \rightarrow e^+e^-$ decays, electron candidates are combined with photon candidates in order to recover some of the energy lost through bremsstrahlung. In the χ_{c1} reconstruction, J/ψ candidates are selected as described above. The associated γ has to satisfy $\text{LAT} <$

0.8, $A_{42} < 0.15$ and has to have an energy greater than 0.15 GeV. The χ_{c1} candidates are required to satisfy $0.35 < M_{\ell+\ell-\gamma} - M_{\ell+\ell-} < 0.45$ GeV/ c^2 , where ℓ represents an electron or a muon. The $\pi^0 \rightarrow \gamma\gamma$ candidates are required to satisfy $0.113 < M_{\gamma\gamma} < 0.153$ GeV/ c^2 . Both photons have to satisfy $\text{LAT} < 0.8$. The energy of the soft photon has to be greater than 0.050 GeV and the energy of the hard photon has to be greater than 0.150 GeV. The $K_s^0 \rightarrow \pi^+ \pi^-$ candidates are required to satisfy $0.489 < M_{\pi^+\pi^-} < 0.507$ GeV/ c^2 . In addition, the K_s^0 flight distance defined as the distance between the reconstructed B and K_s^0 vertices must exceed 1 mm, and the angle between the K_s^0 momentum and its flight direction in the plane transverse to the beam axis must be less than 0.2 rad. The K^{*0} and K^{*+} candidates are required to satisfy $0.796 < M_{K\pi} < 0.996$ GeV/ c^2 and $0.792 < M_{K\pi} < 0.992$ GeV/ c^2 , respectively. In addition, for the sake of suppressing background from events with soft pions, for channels having a π^0 in the final state, the cosine of the angle between the K momentum and the B momentum in the K^* rest frame has to be less than 0.8.

The B candidates are reconstructed by combining charmonium and kaon candidates and are characterized by two kinematic variables: the difference between the reconstructed energy of the B candidate and the beam energy in the center-of-mass frame $\Delta E = E_B^* - E_{beam}^*$, and the beam energy-substituted mass m_{ES} , defined as $m_{ES} \equiv \sqrt{E_{beam}^{*2} - \mathbf{p}_B^{*2}}$, where the $*$ refers to quantities in the center-of-mass and \mathbf{p}_B is the B momentum. For a correctly reconstructed B meson, ΔE is expected to peak at zero and the energy-substituted mass m_{ES} at the B meson mass, 5.279 GeV/ c^2 . Only one reconstructed B meson is allowed per event. For events that have multiple candidates, the candidate having the smallest $|\Delta E|$ is chosen. The analysis is performed in a region of the m_{ES} vs ΔE plane defined by $5.2 < m_{ES} < 5.3$ GeV/ c^2 and $-0.12 < \Delta E < 0.12$ GeV. A ΔE channel-dependent signal region is subsequently defined. The m_{ES} distributions within the ΔE signal region for candidate events are shown on Figure 1.

There are two components to the residual background in the ΔE signal region: the combinatorial background and a peaking component (component of the background that peaks at the same values of ΔE and m_{ES} as the signal). The number of signal events N_S is determined from the number of candidate events, N_{cand} , after subtracting the peaking background. For this purpose, the m_{ES} distribution within the ΔE signal region is fitted to the sum of an ARGUS function [7], which models the combinatorial background, and a Gaussian function. The value of N_{cand} is given by the integral of the Gaussian component. There are two contributions to the peaking background. The first is the cross-feed component that is due to $B \rightarrow (c\bar{c}) K^*$ events from the three channels other than the one under consideration. The second contribution is from other B decays with a J/ψ or $\psi(2S)$ in the final

state. To determine the extent of peaking background, the m_{ES} distribution for simulated $B\bar{B}$ events is fitted within the ΔE signal region by an ARGUS function and a Gaussian function. The peaking background is taken as the integral of the Gaussian portion.

The branching fractions are obtained as:

$$BF \equiv \frac{N_S}{N_{B\bar{B}} \times \epsilon \times f}, \quad (1)$$

where $N_{B\bar{B}}$ is the number of $B\bar{B}$ events, ϵ is the selection efficiency and f is the total secondary branching fraction. For channels with a K^* in the final state, the cross-feed contribution depends on the branching fractions that are being measured. It was estimated by an iterative procedure and found to be small.

The systematic errors arise from the uncertainty on the number of $B\bar{B}$ events (1.1%), the secondary branching fractions (taken from Ref. [8]), the estimate of the selection efficiency, and the knowledge of the background. For the tracking efficiency, an error of 1.3% per track has been used. For the particle identification efficiency, the systematic error varies between 0.2% and 3.7%. The uncertainty on the detection and energy measurement of photons is 2.5%, common to all channels, plus a small channel-dependent correction. The uncertainty on the π^0 reconstruction is 5.0% for all channels, plus a channel-dependent correction. The overall selection efficiency depends on the angular distribution. The efficiency can be written as $\epsilon = a + A_0 b$, where a and b are obtained from the K^* helicity angle by $a = 3/4 \int (1 - \cos^2 \theta_{K^*}) \epsilon(\theta_{K^*}) \sin(\theta_{K^*}) d\theta_{K^*}$ and $b = 3/4 \int (3 \cos^2 \theta_{K^*} - 1) \epsilon(\theta_{K^*}) \sin(\theta_{K^*}) d\theta_{K^*}$, and A_0 is the fraction of the longitudinal K^* polarization [9]. The values of a and b are obtained from simulation. A systematic error varying from 3.4% to 8.6% has been derived for the branching fractions. In the default fit, the shape parameter of the ARGUS function is not constrained. To determine a systematic error due to the combinatorial background, a second fit with the shape parameter of the ARGUS function fixed to the value obtained from fitting the data in the ΔE sideband region was performed. The systematic uncertainty on the combinatorial background has been taken as 50% of the difference between the number of events obtained from the default fit and from the second fit. For the cross-feed component to the peaking background, the uncertainty of the corresponding branching fractions, taken from Ref. [8], has been assigned as the systematic error. For the contribution coming from other B decays with a J/ψ or a $\psi(2S)$ in the final state, a 50% error has been assigned, accounting for the poor knowledge of the branching fractions of the contributing decay modes. Overall, the dominant contribution to the systematic error is from the secondary-decay branching fractions in the case of $\psi(2S) K^{(*)}$ and $\chi_{c1} K$, the efficiency determination in the case of J/ψ




FIG. 1: m_{ES} distributions and fits within the ΔE signal region. From left to right, the columns show the distributions for the J/ψ , $\psi(2S)$ and χ_{c1} channels. From top to bottom, the rows show the distributions for the $B \rightarrow (c\bar{c}) K_S^0$ ($K_S^0 \pi^0$), $B \rightarrow (c\bar{c}) K^{*0}$ ($K^+ \pi^-$), $B \rightarrow (c\bar{c}) K^{*+}$ ($K_S^0 \pi^+$), $B \rightarrow (c\bar{c}) K^{*+}$ ($K^+ \pi^0$), $B \rightarrow (c\bar{c}) K_S^0$ and $B \rightarrow (c\bar{c}) K^+$ decay modes. The dashed and dotted lines show the combinatorial and peaking backgrounds, respectively.

$K^{(*)}$, and the background subtraction in the case of $\chi_{c1} K^*$.

The branching fraction measurements are summarized in Table II. We have taken $R^{+/0}$, the ratio $\Gamma(\Upsilon(4S) \rightarrow B^+ B^-) / \Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0)$, to be unity for the determination of the branching fractions. Assuming isospin invariance in the $B \rightarrow \text{charmonium } K (K^*)$ decays, we

TABLE II: Measured branching fractions for exclusive decays of B mesons to charmonium and kaon final states. The first error is statistical and the second systematic.

Channel	Branching fraction ($\times 10^{-4}$)
$B^0 \rightarrow J/\psi K^{*0}$	$13.09 \pm 0.26 \pm 0.77$
$B^+ \rightarrow J/\psi K^{*+}$	$14.54 \pm 0.47 \pm 0.97$
$B^+ \rightarrow J/\psi K^+$	$10.61 \pm 0.15 \pm 0.48$
$B^0 \rightarrow J/\psi K^0$	$8.69 \pm 0.22 \pm 0.30$
$B^0 \rightarrow \psi(2S) K^{*0}$	$6.49 \pm 0.59 \pm 0.97$
$B^+ \rightarrow \psi(2S) K^{*+}$	$5.92 \pm 0.85 \pm 0.89$
$B^+ \rightarrow \psi(2S) K^+$	$6.17 \pm 0.32 \pm 0.44$
$B^0 \rightarrow \psi(2S) K^0$	$6.46 \pm 0.65 \pm 0.51$
$B^0 \rightarrow \chi_{c1} K^{*0}$	$3.27 \pm 0.42 \pm 0.64$
$B^+ \rightarrow \chi_{c1} K^{*+}$	$2.94 \pm 0.95 \pm 0.98$
$B^+ \rightarrow \chi_{c1} K^+$	$5.79 \pm 0.26 \pm 0.65$
$B^0 \rightarrow \chi_{c1} K^0$	$4.53 \pm 0.41 \pm 0.51$

can compute $R^{+/0}$. Using the ratio of the charged to neutral B meson lifetimes $\tau_{B^+} / \tau_{B^0} = 1.086 \pm 0.017$ [8], we obtain:

$$R^{+/0} \equiv \frac{\Gamma(\Upsilon(4S) \rightarrow B^+ B^-)}{\Gamma(\Upsilon(4S) \rightarrow B^0 \bar{B}^0)} = 1.06 \pm 0.02 \pm 0.03, \quad (2)$$

where the first error is statistical and the second systematic. The ratios of the branching fractions for $B \rightarrow (c\bar{c}) K^*$ and $B \rightarrow (c\bar{c}) K$ for the three $(c\bar{c})$ states are presented in Table III. For each of the charmonium states, the average of the charged and neutral measurements is also shown.

TABLE III: Results for ratios of the branching fractions for charmonium and a K^* versus charmonium and a K . The first error is statistical and the second systematic.

Ratio	Result
$\mathcal{B}(B^0 \rightarrow J/\psi K^{*0}) / \mathcal{B}(B^0 \rightarrow J/\psi K^0)$	$1.51 \pm 0.05 \pm 0.08$
$\mathcal{B}(B^+ \rightarrow J/\psi K^{*+}) / \mathcal{B}(B^+ \rightarrow J/\psi K^+)$	$1.37 \pm 0.05 \pm 0.08$
$\mathcal{B}(B \rightarrow J/\psi K^*) / \mathcal{B}(B \rightarrow J/\psi K)$	$1.44 \pm 0.04 \pm 0.06$
$\mathcal{B}(B^0 \rightarrow \psi(2S) K^{*0}) / \mathcal{B}(B^0 \rightarrow \psi(2S) K^0)$	$1.00 \pm 0.14 \pm 0.09$
$\mathcal{B}(B^+ \rightarrow \psi(2S) K^{*+}) / \mathcal{B}(B^+ \rightarrow \psi(2S) K^+)$	$0.96 \pm 0.15 \pm 0.09$
$\mathcal{B}(B \rightarrow \psi(2S) K^*) / \mathcal{B}(B \rightarrow \psi(2S) K)$	$0.98 \pm 0.10 \pm 0.07$
$\mathcal{B}(B^0 \rightarrow \chi_{c1} K^{*0}) / \mathcal{B}(B^0 \rightarrow \chi_{c1} K^0)$	$0.72 \pm 0.11 \pm 0.12$
$\mathcal{B}(B^+ \rightarrow \chi_{c1} K^{*+}) / \mathcal{B}(B^+ \rightarrow \chi_{c1} K^+)$	$0.51 \pm 0.17 \pm 0.16$
$\mathcal{B}(B \rightarrow \chi_{c1} K^*) / \mathcal{B}(B \rightarrow \chi_{c1} K)$	$0.65 \pm 0.09 \pm 0.10$

Finally, we have measured the charge asymmetries

$$A \equiv \frac{\mathcal{B}(B^+ \rightarrow (c\bar{c}) K^{+(*)}) - \mathcal{B}(B^- \rightarrow (c\bar{c}) K^{-(*)})}{\mathcal{B}(B^+ \rightarrow (c\bar{c}) K^{+(*)}) + \mathcal{B}(B^- \rightarrow (c\bar{c}) K^{-(*)})}, \quad (3)$$

using efficiencies determined separately for the two charges. The results are presented in Table IV. No statistically significant asymmetry is observed.

TABLE IV: Results for charge asymmetries. The first error is statistical and the second systematic.

Final state	Asymmetry
$J/\psi K^+$	$-0.030 \pm 0.014 \pm 0.010$
$J/\psi K^{*+}$	$0.048 \pm 0.029 \pm 0.016$
$\psi(2S) K^+$	$0.052 \pm 0.059 \pm 0.020$
$\psi(2S) K^{*+}$	$-0.077 \pm 0.207 \pm 0.051$
$\chi_{c1} K^+$	$0.003 \pm 0.076 \pm 0.017$
$\chi_{c1} K^{*+}$	$-0.471 \pm 0.378 \pm 0.268$

In summary, branching fraction measurements of exclusive B decays to charmonium (J/ψ , $\psi(2S)$ and χ_{c1}) and K or K^* have been presented. Our results for J/ψ and $\psi(2S)$ are in good agreement with previous measurements [8] and exhibit comparable or superior precision. Our χ_{c1} results have much better precision than earlier measurements. The $B^+ \rightarrow \chi_{c1} K^{*+}$ mode was previously unmeasured. Assuming isospin invariance, we find the ratio of charged- to neutral- B meson production on the $\Upsilon(4S)$ resonance to be compatible with unity within 1.7 standard deviations. No direct CP violation has been observed in the charge asymmetries.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and

NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Also with Università della Basilicata, Potenza, Italy

† Deceased

- [1] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D. **65**, 032001 (2002).
- [2] T. Brown, S. Pakvasa and S.F. Tuan, Phys. Lett. **B136**, 117 (1984); I. Dunietz, Phys. Lett. **B316**, 561 (1993).
- [3] *BELLE* Collaboration, K. Abe *et al.*, Phys. Rev. D. **67**, 032003 (2003).
- [4] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instr. and Methods **A 479**, 1 (2002).
- [5] A. Drescher *et al.*, Nucl. Instrum. Methods **A237**, 464 (1985).
- [6] R. Sinkus and T. Voss, Nucl. Instrum. Methods **A391**, 360 (1997).
- [7] *ARGUS* Collaboration, H. Albrecht *et al.*, Z. Phys. C **48**, 543 (1990).
- [8] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. **B592**, 1 (2004).
- [9] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 241801 (2001).